

WEST Search History

DATE: Wednesday, November 09, 2005

<u>Hide?</u>	<u>Set Name</u>	<u>Query</u>	<u>Hit Count</u>
<i>DB=PGPB,USPT,USOC,EPAB,JPAB,DWPI; PLUR=YES; OP=ADJ</i>			
<input type="checkbox"/>	L6	L5 and apoe4 and transgenic	42
<input type="checkbox"/>	L5	435/455,463,320.1,325.ccls.	35986
<input type="checkbox"/>	L4	L3 and apoe4	13
<input type="checkbox"/>	L3	800/18,3,9,13,13,21,22.ccls.	1798
<input type="checkbox"/>	L2	apoe4 with transgenic	40
<input type="checkbox"/>	L1	apoe4 near transgenic	8

END OF SEARCH HISTORY

(FILE 'HOME' ENTERED AT 16:24:28 ON 09 NOV 2005)

FILE 'CAPLUS, MEDLINE, EMBASE, BIOSIS, LIFESCI' ENTERED AT 16:24:36 ON 09 NOV 2005

L1 85 S APOE4 WITH TRANSGENIC
L2 11 S L1 AND (THR OR ARG OR THREONINE OR ARGININE)
L3 4 DUP REM L2 (7 DUPLICATES REMOVED)
L4 4 S L1 AND 61
L5 1 DUP REM L4 (3 DUPLICATES REMOVED)
L6 78 S L1 AND HUMAN
L7 31 S L6 AND PY<=2001
L8 14 DUP REM L7 (17 DUPLICATES REMOVED)
L9 938 S WEISGRABER K?/AU
L10 11 S L1 AND L2
L11 0 S L9 AND L1
L12 205 S L9 AND APOE4
L13 12 S L12 AND TRANSGEN?
L14 4 DUP REM L13 (8 DUPLICATES REMOVED)
L15 1217 S FARESE R?/AU
L16 8 S L15 AND APOE4
L17 4 DUP REM L16 (4 DUPLICATES REMOVED)
L18 39 S RAFFAI RO?/AU
L19 13 S L18 AND APOE4
L20 6 DUP REM L19 (7 DUPLICATES REMOVED)
L21 1900 S DONG LI?/AU
L22 21 S L21 AND APOE4
L23 10 DUP REM L22 (11 DUPLICATES REMOVED)



All Databases

PubMed

Nucleotide

Protein

Genome

Structure

PMC

Taxonomy

Books

C

Search Gene

for

Go

Clear

current records

Limits Preview/Index History Clipboard Details

Display Full Report

Show 20

Send to

All: 1 Genes Genomes: 1 SNP GeneView: 1


1: Apoe apolipoprotein E [Mus musculus]

GeneID: 11816 Locus tag: MGI:88057

updated 06-Nov-2005

Summary**Official Symbol:** Apoe and **Name:** apolipoprotein E provided by [Mouse Genome Informatics](#)**Gene type:** protein coding**Gene name:** Apoe**Gene description:** apolipoprotein E**RefSeq status:** Provisional**Organism:** *Mus musculus***Lineage:** Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Sciurognathi; Muroidea; Muridae; Murinae; *Mus***Gene aliases:** AI255918**Genomic regions, transcripts, and products**

(minus strand) RefSeq below

[NC_000073](#)**Genomic context**[See Apoe in MapViewer](#)

chromosome: 7; Locations: 7 A2; 7 4.0 cM

16547339 ►

16618714 ►

**Bibliography**Gene References into Function (GeneRIF): [Submit](#)

?

[PubMed links](#)**GeneRIFs:**

1. Scarb1/apoE knockout mice have coronary heart disease not involving lymphocytes [PubMed](#)
2. ApoE mediates the presentation of serum-borne lipid antigens and can be secreted by antigen presenting cells as a mechanism to survey the local environment to capture antigens or to transfer microbial lipids from infected cells to bystander APCs [PubMed](#)

Entrez Gene**Table Of Content**

Summary
 Genomic regions, tran
 Genomic context
 Bibliography
 Alleles
 General gene inform
 General protein inform
 Reference Sequences
 Related Sequences
 Additional Links

Links

MGC cDNA clone
 Conserved Domains
 Genome
 GEO Profiles
 HomoloGene
 Map Viewer
 Nucleotide
 Full text in PMC
 Probe
 Protein
 PubMed
 PubMed (GeneRIF)
 SNP
 SNP: Genotype
 SNP: GeneView
 Taxonomy
 UniSTS
 Ensembl
 Evidence Viewer
 Gene Expression Data
 KEGG
 MGC
 MGI
 ModelMaker
 UCSC
 UniGene
 LinkOut

Entrez Gene In[Feedback](#)[Subscription](#)

3. ApoE4 mice develop a constellation of changes that mimic the pathology associated with human age-related macular degeneration [PubMed](#)

4. protein engineering studies show that a destabilized conformation promotes apoE4 lipid binding [PubMed](#)

5. additive effects of apoE and clusterin on influencing amyloid-beta deposition [PubMed](#)

6. A cholesterol-independent role of apolipoprotein E in atherosclerosis regression is critical for lipid removal from the fibrotic component of the plaque but not from the foam cell-rich layer beneath the endothelium. [PubMed](#)

7. Human CRP transgene expression is thus up-regulated in apoE-deficient mice, apparently reflecting altered estrogen levels, despite the absence of other systemic signs of inflammation [PubMed](#)

8. In addition to the effect of apoE on lipid trafficking, apoE may also influence the astroglial response to damage [PubMed](#)

9. Chronic renal failure aggravates atherosclerosis in apoE(-/-) mice. [PubMed](#)

10. ApoE has a physiologic role as a regulator of osteoblast function [PubMed](#)

11. Effects of a standardized freeze-dried powder made from fresh grapes on the development of atherosclerotic lesions in apolipoprotein E deficient mice. [PubMed](#)

12. Increased atherosclerosis is accompanied by increases in VCAM-1 and P-selectin levels in the two apoE(-/-) mouse strains, the high HDL level may protect against atherosclerosis by inhibiting the expression of adhesion molecules in BALB/c.apoE(-/-) mice. [PubMed](#)

13. autoregulation of the LX α gene facilitates induction of apoE in mouse adipose tissue [PubMed](#)

14. important in HSV-1 colonization of ovaries [PubMed](#)

15. In a mouse model of restenosis, a subphysiological level of apoE was associated with beneficial effects on lesion size/composition. ApoE has a generally important protective function in the arterial wall. [PubMed](#)

16. Results indicate that adipose tissue expression of apolipoprotein E is modulated by physiologic regulators of insulin sensitivity. [PubMed](#)

17. Improvement in insulin resistance induced by a high-fat diet, in apoE knockout mice of vasculopathy. did not alter plaque composition, [PubMed](#)

18. activation of extracellular signal-regulated kinase has a role in increasing tau phosphorylation in apolipoprotein E4 transgenic mice [PubMed](#)

19. Deficiency of apoE protein in macrophages attenuates the ingestion of apoptotic cells in vitro and results in impaired clearance of apoptotic cell remnants and a functionally relevant systemic proinflammatory condition in vivo. [PubMed](#)

20. Intravital microscopy of the common carotid artery revealed a significantly greater number of leukocytes rolling on the vessel walls in apoE-/-pGI receptor-/- mice. [PubMed](#)

21. The apoE4 isoform of apolipoprotein E (apoE), which is the major genetic risk factor of Alzheimer disease. [PubMed](#)

22. plays a significant role in host defense against candidiasis. [PubMed](#)

23. Review. This review will examine our present understanding of the pathology and progression of plaques in Apo-E-deficient mice and highlight some of the nutritional, pharmacological, and genetic studies that have enhanced this understanding. [PubMed](#)

24. ApoE3 has a role in liver expression of fatty acid binding protein, as shown in transgenic mice models [PubMed](#)

25. apoE metabolism requires ABCA1 protein in the central nervous system [PubMed](#)

26. Results describe the relative roles of apolipoprotein E isoforms in low density lipoprotein receptor (LDLR)- and non-LDLR-mediated very low density lipoprotein (VLDL) clearance. [PubMed](#)

27. The accelerated renal injury that was observed in diabetic apo E-KO mice was attenuated by approaches that inhibit renal AGE accumulation. [PubMed](#)

28. serum chemokine levels are potential markers for atherosclerosis susceptibility in C3H and C57BL apoE(-/-) mice fed on a normal rodent diet [PubMed](#)

29. uremia markedly accelerates atherogenesis in apolipoprotein E-deficient mice. [PubMed](#)

30. Results show that alterations in DNA methylation profiles are early markers of atherosclerosis in an apolipoprotein E-null mouse model and may play a causative role in atherogenesis. [PubMed](#)

31. Knockout mice are at risk for cognitive deficits. [PubMed](#)

32. Apolipoprotein E promotes astrocyte colocalization and degradation of deposited amyloid-beta peptides. [PubMed](#)

33. After sirolimus treatment, the cholesterol content of the aortic arch was reduced in apoe knockout mice compared to normal controls. [PubMed](#)

34. estradiol facilitates neurite growth through an ApoE-dependent mechanism. [PubMed](#)

35. Low plasma apoE (1-3 x 10(-8) M) suppresses atherosclerosis by as yet undefined mechanisms, not dependent on the presence of apoA-I or HDL or an increased capacity of serum acceptors for Free Cholesterol efflux [PubMed](#)

36. ApoE deficient mice had significantly increased lymphocyte proliferation responses to both myelin antigens and mitogens and significantly more infiltrating lesions in the central nervous system (CNS) in histopathology [PubMed](#)

37. generation and characterization of two transgenic mouse lines expressing human ApoE2 in neurons and glial cells [PubMed](#)

38. use of a model of conditional gene repair to examine metabolism [PubMed](#)

39. Conditional disruption of the peroxisome proliferator-activated receptor gamma gene in mice results in lowered expression of ABCA1, ABCG1, and apoE in macrophages and reduced cholesterol efflux. [PubMed](#)

40. modulating transport of polyunsaturated phospholipid molecular species in synaptic plasma membranes. [PubMed](#)

41. Results identify the cyclooxygenase-2 gene as a target of APOE signaling, link HDL and APOE to prostacyclin receptor IP action, and describe a potential new basis for the cardioprotective effect of HDL and APOE. [PubMed](#)

42. ApoE normally associates with acetylcholinesterase in the synaptic cleft of slow muscles, modulating the activity of the enzyme and therefore quantal size. [PubMed](#)

43. neuronal expression of apoE is regulated by a diffusible factor or factors released from astrocytes [PubMed](#)

44. impact of telomere attrition on atherogenesis induced by dietary cholesterol in apolipoprotein E (apoE)-deficient mice [PubMed](#)

45. In aged female apoE0/0 mice water maze performance was impaired with search strategies. In parallel, increased corticosterone concentrations were measured in apoE0/0 mice in response to novelty and during the circadian cycle. [PubMed](#)

46. Lack of both NPRA and apoE synergistically enhances cardiac hypertrophy, suggesting roles for Npr1 as well as Apoe in regulation of hypertrophic cell growth. [PubMed](#)

47. Results demonstrate that apoE4-202 functions even in the presence of clearance-defective apoE2. [PubMed](#)

48. isoform-specific role in mediating the systemic and brain inflammatory responses [PubMed](#)

49. TR4 can also regulate apolipoprotein E, C-I, and C-II gene expression via the TR4 response element within the hepatic control region [PubMed](#)

50. Brain regional differences in the production of apoE throughout the estrous cycle were noted. Exogenous estradiol has regionally specific effects on apoE expression. Regional variability appears to vary as a function of the estrogen receptor subtype [PubMed](#)

51. role in Alzheimer's disease-like neurodegeneration and behavioral deficits in transgenic mice [PubMed](#)

52. apoE inhibits cell migration via cAMP-dependent protein kinase A activation as a consequence of its binding to LRP-1 [PubMed](#)

53. After adoptive regulatory T cells type 1 transfer in female apolipoprotein E-knockout mice, developing atherosclerotic plaques had fewer macrophages and T cells than plaques of control mice	PubMed
54. apoE deficiency in apoE knockout mice leads to a deficit in olfactory function, suggesting an important role for apoE in the olfactory system	PubMed
55. ApoE(-/-)/IL-4(-/-) mice showed a 58% and 64% decrease in disease in aortic arch compared to apoE(-/-) and apoE(-/-)/IL-12(-/-) mice, respectively, and a 78% decrease in thoracic lesions compared to apoE(-/-)/IL-12(-/-).	PubMed
56. Apolipoprotein E facilitates cerebral amyloid angiopathy and spontaneous hemorrhage in amyloid precursor protein transgenic mice.	PubMed
57. ApoB-48 and apoB-100 differentially influence the expression of type-III hyperlipoproteinemia in APOE*2 mice	PubMed
58. adrenal gland cholesterol metabolism in apolipoprotein E deficient mice	PubMed
59. lack of lipoprotein receptor-related protein 5 and apolipoprotein E is seen in severe hypercholesterolemia, impaired fat tolerance, and advanced atherosclerosis	PubMed
60. recycling of apolipoprotein E in primary cultures of mouse hepatocytes	PubMed
61. Following complete spinal cord transection Apo-E is upregulated in neutrophils and macrophages at the injury site and is found at later times in astrocytes during remodeling of white matter tracts, notably in degenerating parts of the fasciculus gracilis.	PubMed
62. Apolipoprotein E deficiency promotes increased oxidative stress and compensatory increases in antioxidants in brain tissue.	PubMed
63. The increase in apoE expression closely correlated in time and spatial distribution with axonal and neuronal degeneration, consistent with a role as an 'injury-response' protein.	PubMed
64. Transgenic mice deficient in apoE display altered levels of mature amyloid precursor protein (APP) as well as increased amounts of both beta-cleaved C-terminal fragments and Abeta peptides.	PubMed
65. Mice lacking apoE appear not to be impaired in spatial memory per se but are deficient in a procedural component of the Morris water maze.	PubMed
66. The effect of angiotensin II on arterial wall stiffness in apolipoprotein E knockout mice.	PubMed
67. involvement of ApoE in the hematogenous route of HSV-1 to the CNS	PubMed
68. Peroxisome proliferator-activated receptor alpha,gamma coagonist LY465608 inhibits macrophage activation and atherosclerosis in apolipoprotein E knockout mice	PubMed
69. in null mice, blockade of platelet-derived growth factor or its receptors transiently delays but does not prevent fibrous cap formation	PubMed
70. The data indicate a neuroprotective role for estrogen in global ischemia, the mechanism of which is apoE-dependent	PubMed
71. regulated expression of gene cluster in macrophages	PubMed
72. regulation of macrophage ApoE expression and processing by extracellular matrix	PubMed
73. helps maintain blood brain barrier; of particular importance after brain injury	PubMed
74. Apolipoprotein E4 potentiates amyloid beta peptide-induced lysosomal leakage and apoptosis in neuronal cells	PubMed
75. Reconstituted discoidal ApoE-phospholipid particles are ligands for the scavenger receptor BI	PubMed
76. ApoE affects the age of onset of Abeta deposition in amyloid precursor protein transgenic mice as well as the level, structure and anatomic distribution of brain Abeta deposits.	PubMed
77. Apo-E knockout middle cerebral artery occlusion mice showed a worsened deficit in locomotor activity, which was significantly correlated with exacerbated cortical lesion volume	PubMed

Alleles



The following allele types are documented at Mouse Genome Informatics ([MGI](#))

- Targeted (knock-in) (1) [PubMed](#)
- Targeted (knock-out) (3) [PubMed](#)

General gene information



Markers

Apoe(e-PCR) (Links: [UniSTS: 141112](#))

Alternate name: MGI:1205863

AI255918(e-PCR) (Links: [UniSTS: 179533](#))

Alternate name: 392587

PMC186328P2(e-PCR) (Links: [UniSTS: 271700](#))

GeneOntology

Provided by [MGI](#)

Function	Evidence	
heparin binding	IEA	
lipid binding	IEA	
lipid transporter activity	IEA	
lipoprotein binding	IDA	PubMed
Process		
cholesterol homeostasis	IMP	PubMed
cholesterol metabolism	IMP	PubMed
lipid transport	IEA	
lipoprotein metabolism	IEA	
response to oxidative stress	IMP	PubMed
transport	IEA	
Component		
chylomicron	IEA	
extracellular region	IEA	
extracellular space	RCA	PubMed

Homology:

Human, Rat

[Map Viewer](#)

Pathways

KEGG pathway: Alzheimer's disease [05010](#)

KEGG pathway: Neurodegenerative Disorders [01510](#)

General protein information



Name: apolipoprotein E

NCBI Reference Sequences (RefSeq)



mRNA Sequence [NM_009696](#)

Source Sequence [BC028816](#)

Product [NP_033826](#) apolipoprotein E

Conserved Domains (1) [summary](#)

[pfam01442: Apolipoprotein; Apolipoprotein A1/A4/E family](#)

Related Sequences



Nucleotide	Protein	Strain
Genomic D00466	BAA00361	BALB/c
mRNA AK010261	None	C57BL/6J
mRNA AK019319	None	C57BL/6J
mRNA AK075843	BAC36000	C57BL/6J
mRNA AK131624	None	C57BL/6J
mRNA AK134921	BAE22338	C57BL/6J
mRNA AK148747	None	C57BL/6J
mRNA AK149111	BAE28740	C57BL/6J
mRNA AK149568	BAE28964	C57BL/6J
mRNA AK150834	BAE29894	C57BL/6J
mRNA AK159105	BAE34821	C57BL/6J
mRNA AK159424	BAE35071	C57BL/6J
mRNA AK159517	BAE35147	C57BL/6J
mRNA BC028816	AAH28816	FVB/N
mRNA BC044785	None	FVB/N
mRNA BC083351	AAH83351	C57BL/6
mRNA CT010212	CAJ18420	
mRNA CT010356	CAJ18564	
mRNA M12414	AAA37251	
mRNA M73490	AAA37252	C57BL/6J
None	P08226	

Additional Links

UniGene [Mm.305152](#)Gene Expression Database (GXD) at MGI [MGI:88057](#)Display [Full Report](#)
 Show 20 Send to

[Restrictions on Use](#) | [Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)



Search for



in these sections

All sections
Gene symbols/names
Accession IDs
Phenotype/Human Disease
Gene Expression
Gene Ontology
Anatomical Dictionary
Phenotype Ontology (MP)

[Advanced search for...](#)



Search Categories

[All Search Tools](#)

[Genes/Markers](#)

[Phenotypes/Alleles](#) (NEW)

[Strains/Polymorphisms](#) (NEW)

[Expression](#)

[Sequences](#)

[Comparative Maps/Data](#)

[Mouse Maps/Data](#)

[Mouse Tumor Biology](#)

[Probes/Clones](#)

[References](#)

[Vocabulary Browsers](#)

[Anatomical Dictionary](#)

[Gene Ontology \(GO\)](#)

[Human Disease](#)

[\(OMIM\)](#) (NEW)

[Phenotype Ontology \(MP\)](#)

[Protein Superfamily](#) (NEW)

[MouseBLAST](#)

[Mouse GBrowse](#)

[IMSR \(Find Mice\)](#)

Tools and Links

[Citing These Resources](#)

[Funding Information](#)

[Warranty Disclaimer](#)

[& Copyright Notice](#)

Send questions and
comments to [User Support](#).

Other Genome Feature Detail

Your Input Welcome

Symbol Name ID	Tg(GFAP-APOE4) 1Hol	Nomenclature
	transgene insertion 1, David M Holtzman MGI:3057183	History
Genetic Map	Chromosome Unknown	
Phenotypes	All phenotypic alleles(1) : Transgenic(1)	
References	J:93487 Sun Y et al., "Glial fibrillary acidic protein-apolipoprotein E (apoE) transgenic mice: astrocyte-specific expression and differing biological effects of astrocyte-secreted apoE3 and apoE4 lipoproteins." J Neurosci 1998 May 1;18(9):3261-72 All references(1)	



last database update
11/07/2005
MGI 3.4